




PCT/EP200 5 / 0 0 0 5 4 2



INVESTOR IN PEOPLE

The Patent Office  
Concept House  
Cardiff Road  
Newport  
South Wales  
NP10 8QQ



## PRIORITY DOCUMENT


SUBMITTED OR TRANSMITTED IN  
COMPLIANCE WITH RULE 17.1(a) OR (b)

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.



Signed

*Anders Grey*

Dated

4 November 2004

## Patents Form 1/77

Patents Act 1977  
(Rule 16)THE PATENT OFFICE  
A

21 JAN 2004

The  
Patent  
Office

1/77

## Request for grant of a patent

(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)

The Patent Office

Cardiff Road  
Newport  
Gwent NP10 8QQ

1.	Your reference	4-33598P1/HO 96		
2.	Patent application number (The Patent Office will fill in this part)	0401336.3		
3.	Full name, address and postcode of the or of each applicant (underline all surnames)	NOVARTIS AG LICHTSTRASSE 35 4056 BASEL SWITZERLAND 7125487005 SWITZERLAND		
	Patent ADP number (if you know it) If the applicant is a corporate body, give the country/state of its incorporation			
4.	Title of invention	Organic Compounds		
5.	Name of your agent (If you have one) "Address for service" in the United Kingdom to which all correspondence should be sent (including the postcode)	Bernard Marsh Novartis Pharmaceuticals UK Limited Patents and Trademarks Wimblehurst Road Horsham West Sussex RH12 5AB 7181522002		
	Patents ADP number (if you know it)			
6.	If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or of each of these earlier applications and (if you know it) the or each application number	Country	Priority application number (if you know it)	Date of filing (day/month/year)
7.	If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application	Number of earlier application	Date of filing (day/month/year)	
8.	Is a statement of inventorship and of right to grant of a patent required in support of this request? (Answer 'Yes' if: a) any applicant named in part 3 is not an inventor, or b) there is an inventor who is not named as an applicant, or c) any named applicant is a corporate body. (see note (d))	Yes		

## Patents Form 1/77

9. Enter the number of sheets for any of the following items you are filing with this form. Do not count copies of the same document

Continuation sheets of this form

Description 19 ✓

Claim(s) 4 ✓

Abstract

Drawing(s)

10. If you are also filing any of the following, state how many against each item.

Priority documents

Translations of priority documents

Statement of inventorship and right to grant of a patent (*Patents Form 7/77*)

Request for preliminary examination and search (*Patents Form 9/77*) 1

Request for substantive examination (*Patents Form 10/77*)


Any other documents  
(please specify)

11.

I/We request the grant of a patent on the basis of this application

Signature

Date



Bernard Marsh

21 January 2004

12. Name and daytime telephone number of person to contact in the United Kingdom

Mr. T. Drew  
(01403) 32 3069

### Warning

*After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.*


### Notes

- If you need help to fill in this form or you have any questions, please contact the Patent Office on 0645 500505.
- Write your answers in capital letters using black ink or you may type them.
- If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.
- Once you have filled in the form you must remember to sign and date it.
- For details of the fee and ways to pay please contact the Patent Office.

Organic Compounds

This invention relates to organic compounds, their preparation and their use as pharmaceuticals.

In one aspect, the present invention provides compounds of formula



in free or salt form, where

Ar is a C<sub>6</sub>-C<sub>15</sub> monovalent aromatic group,

R<sup>1</sup> is hydrogen, phenyl optionally substituted by one or more substituents selected from halogen, cyano, hydroxy, C<sub>1</sub>-C<sub>8</sub>-alkyl, C<sub>1</sub>-C<sub>8</sub>-haloalkyl, C<sub>1</sub>-C<sub>8</sub>-alkoxy, C<sub>1</sub>-C<sub>8</sub>-alkoxy-C<sub>1</sub>-C<sub>8</sub>-alkyl and acyloxy, or R<sup>1</sup> is a 5- or 6- membered monovalent heterocyclic group,

R<sup>2</sup> is hydrogen, C<sub>1</sub>-C<sub>8</sub>-alkyl, acyl or -CON(R<sup>3</sup>)R<sup>4</sup>,

R<sup>3</sup> and R<sup>4</sup> are each independently hydrogen or C<sub>1</sub>-C<sub>8</sub>-alkyl, or together with the nitrogen atom to which they are attached denote a 5- or 6- membered heterocyclic group, and

Y is a pyrimidinyl or pyridazinyl group, optionally substituted by at least one C<sub>1</sub>-C<sub>8</sub>-alkyl, C<sub>1</sub>-C<sub>8</sub>-alkoxy, C<sub>1</sub>-C<sub>8</sub>-alkylthio, C<sub>1</sub>-C<sub>8</sub>-alkylamino, di(C<sub>1</sub>-C<sub>8</sub>-alkyl)amino or acylamino group.

Terms used in the specification have the following meanings :

"C<sub>1</sub>-C<sub>8</sub>-alkyl" as used herein denotes straight chain or branched C<sub>1</sub>-C<sub>8</sub>-alkyl, which may be, for example, methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, straight or branched pentyl, straight or branched hexyl, straight or branched heptyl, or straight or branched octyl. Preferably, C<sub>1</sub>-C<sub>8</sub>-alkyl is C<sub>1</sub>-C<sub>4</sub>-alkyl.

"C<sub>1</sub>-C<sub>8</sub>-alkoxy" as used herein denotes straight chain or branched C<sub>1</sub>-C<sub>8</sub>-alkoxy which may be, for example, methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, isobutoxy, sec-butoxy, tert-butoxy, straight or branched pentoxy, straight or branched hexyloxy, straight or branched heptyloxy, or straight or branched octyloxy. Preferably, C<sub>1</sub>-C<sub>8</sub>-alkoxy is C<sub>1</sub>-C<sub>4</sub>-alkoxy.

"C<sub>1</sub>-C<sub>8</sub>-alkylthio" as used herein denotes straight chain or branched C<sub>1</sub>-C<sub>8</sub>-alkylthio which may be, for example, methylthio, ethylthio, n-propylthio, isopropylthio, n-butylthio, isobutylthio, sec-butylthio, tert-butylthio, straight or branched pentylthio, straight or branched hexylthio, straight or branched heptylthio, or straight or branched octylthio. Preferably, C<sub>1</sub>-C<sub>8</sub>-alkylthio is C<sub>1</sub>-C<sub>4</sub>-alkylthio.

"di(C<sub>1</sub>-C<sub>8</sub>-alkyl)amino" as used herein denotes amino substituted by two C<sub>1</sub>-C<sub>8</sub>-alkyl groups as hereinbefore defined, which may be the same or different. Preferably, di(C<sub>1</sub>-C<sub>8</sub>-alkyl)amino is di(C<sub>1</sub>-C<sub>4</sub>-alkyl)amino.

"C<sub>1</sub>-C<sub>8</sub>-haloalkyl" as used herein denotes C<sub>1</sub>-C<sub>8</sub>-alkyl as hereinbefore defined substituted by one or more halogen atoms, preferably one, two or three halogen atoms, preferably fluorine or chlorine atoms. Preferably C<sub>1</sub>-C<sub>8</sub>-haloalkyl is C<sub>1</sub>-C<sub>4</sub>-alkyl substituted by one, two or three fluorine or chlorine atoms.

"C<sub>1</sub>-C<sub>8</sub>-alkoxy-C<sub>1</sub>-C<sub>8</sub>-alkyl" as used herein denotes C<sub>1</sub>-C<sub>8</sub>-alkyl as hereinbefore defined substituted by C<sub>1</sub>-C<sub>8</sub>-alkoxy as hereinbefore defined.

"C<sub>1</sub>-C<sub>8</sub>-alkylcarbonyl", "C<sub>1</sub>-C<sub>8</sub>-haloalkylcarbonyl" and "C<sub>1</sub>-C<sub>8</sub>-alkoxycarbonyl" as used herein denote C<sub>1</sub>-C<sub>8</sub>-alkyl, C<sub>1</sub>-C<sub>8</sub>-haloalkyl or C<sub>1</sub>-C<sub>8</sub>-alkoxy respectively as hereinbefore defined attached by a carbon atom to a carbonyl group.

"Acyl" as used herein denotes alkylcarbonyl, for example C<sub>1</sub>-C<sub>8</sub>-alkylcarbonyl where C<sub>1</sub>-C<sub>8</sub>-alkyl may be one of the C<sub>1</sub>-C<sub>8</sub>-alkyl groups hereinbefore mentioned, optionally substituted by one or more halogen atoms; cycloalkylcarbonyl, for example C<sub>3</sub>-C<sub>8</sub>-cycloalkylcarbonyl where C<sub>3</sub>-C<sub>8</sub>-cycloalkyl may be, for example, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl or cyclooctyl; 5- or 6- membered heterocyclylcarbonyl having one or more, preferably one or two, hetero atoms selected from nitrogen, oxygen and sulfur in the ring, such as furylcarbonyl or pyridylcarbonyl; arylcarbonyl, for example C<sub>6</sub>-C<sub>10</sub>-arylcarbonyl such as benzoyl; or aralkylcarbonyl, for example C<sub>6</sub> to C<sub>10</sub>-aryl-C<sub>1</sub>-C<sub>4</sub>-alkylcarbonyl such as benzylcarbonyl or phenylethylcarbonyl.

"Acyloxy" as used herein denotes alkylcarbonyloxy, for example C<sub>1</sub>-C<sub>8</sub>-alkylcarbonyloxy where C<sub>1</sub>-C<sub>8</sub>-alkyl may be one of the C<sub>1</sub>-C<sub>8</sub>-alkyl groups hereinbefore mentioned, optionally substituted by one or more halogen atoms; cycloalkylcarbonyloxy, for example C<sub>3</sub>-C<sub>8</sub>-

cycloalkylcarbonyloxy where C<sub>3</sub>-C<sub>8</sub>-cycloalkyl may be, for example, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl or cyclooctyl; 5- or 6- membered heterocyclylcarbonyloxy having one or two hetero atoms selected from nitrogen, oxygen and sulfur in the ring, such as furylcarbonyloxy or pyridylcarbonyloxy; arylcarbonyloxy, for example C<sub>6</sub>-C<sub>10</sub>-arylcarbonyloxy such as benzoyloxy; or alkylcarbonyloxy, for example C<sub>6</sub> to C<sub>10</sub>-aryl-C<sub>1</sub>-C<sub>4</sub>-alkylcarbonyloxy such as benzylcarbonyloxy or phenylethylcarbonyloxy. Preferably acyloxy is C<sub>1</sub>-C<sub>4</sub>-alkylcarbonyloxy.

"Acylamino" as used herein denotes amino substituted by acyl as hereinbefore defined. Preferably it is C<sub>1</sub>-C<sub>4</sub>-alkylcarbonylamino.

"Halogen" as used herein may be fluorine, chlorine, bromine or iodine; preferably it is fluorine or chlorine.

Ar may be, for example, phenyl optionally substituted by one or more substituents, for example, one, two or three substituents selected from halogen, cyano, C<sub>1</sub>-C<sub>8</sub>-alkyl, C<sub>1</sub>-C<sub>8</sub>-haloalkyl or naphthyl. Ar is preferably phenyl optionally substituted by halogen, cyano or C<sub>1</sub>-C<sub>8</sub>-alkyl, preferably meta or para to the indicated thiazole ring.


R<sup>1</sup> may be, for example, hydrogen, phenyl optionally substituted by halogen, cyano, hydroxy, C<sub>1</sub>-C<sub>8</sub>-alkyl, C<sub>1</sub>-C<sub>8</sub>-haloalkyl, C<sub>1</sub>-C<sub>8</sub>-alkoxy, C<sub>1</sub>-C<sub>8</sub>-alkoxy-C<sub>1</sub>-C<sub>8</sub>-alkyl, carboxy, C<sub>1</sub>-C<sub>8</sub>-alkoxycarbonyl or C<sub>1</sub>-C<sub>8</sub>-alkylcarbonyloxy, or a monovalent 5- or 6- membered heterocyclic group having one, two or three ring hetero atoms selected from nitrogen, oxygen and sulfur, such as pyrrolyl, imidazolyl, triazolyl, pyridyl, oxopyridyl, piperidyl, pyridazinyl, pyrimidinyl, pyrazinyl, pyrazolyl, pyrazolinyl, piperazinyl, morpholinyl, furyl, pyranyl, thienyl or thiazolyl, optionally substituted by one or more substituents selected from C<sub>1</sub>-C<sub>8</sub>-alkyl, hydroxy or C<sub>1</sub>-C<sub>8</sub>-alkoxy. Preferably R<sup>1</sup> is phenyl optionally substituted by cyano, carboxy or C<sub>1</sub>-C<sub>4</sub>-alkoxy, or a monovalent 6-membered N-heterocyclic group, especially pyridyl, C<sub>1</sub>-C<sub>4</sub>-alkylpyridyl, C<sub>1</sub>-C<sub>4</sub>-alkoxypyridyl or pyrazinyl.

R<sup>2</sup> may be, for example, hydrogen, C<sub>1</sub>-C<sub>8</sub>-alkyl, formyl, C<sub>1</sub>-C<sub>8</sub>-alkylcarbonyl, C<sub>1</sub>-C<sub>8</sub>-haloalkylcarbonyl, C<sub>3</sub>-C<sub>8</sub>-cycloalkylcarbonyl, phenylcarbonyl in which the phenyl moiety is optionally substituted by halogen, cyano, hydroxy, C<sub>1</sub>-C<sub>8</sub>-alkyl or C<sub>1</sub>-C<sub>8</sub>-alkoxy, heterocyclylcarbonyl in which the heterocyclyl group is 5- or 6-membered and has one or more, preferably one or two, ring hetero atoms selected from nitrogen, oxygen and sulfur, or


a group  $-\text{CON}(\text{R}^3)\text{R}^4$ . Preferably  $\text{R}^2$  is hydrogen,  $\text{C}_1\text{-C}_4$ -alkylcarbonyl, 5-membered heterocyclylcarbonyl, especially furylcarbonyl, or phenylcarbonyl in which the phenyl moiety is optionally substituted by  $\text{C}_1\text{-C}_8$ -alkoxy, especially  $\text{C}_1\text{-C}_4$ -alkoxyphenylcarbonyl.

Where present,  $\text{R}^3$  and  $\text{R}^4$  may each independently be, for example, hydrogen or  $\text{C}_1\text{-C}_4$ -alkyl, or together with the nitrogen atom to which they are attached may denote a 5-membered heterocyclyl group such as pyrrolyl or pyrrolidinyl or a 6-membered heterocyclyl group such as pyridyl, piperidyl, piperazinyl or morpholinyl. Preferably  $\text{R}^3$  and  $\text{R}^4$ , where present, are each  $\text{C}_1\text{-C}_8$ -alkyl, especially methyl, or together with the nitrogen atom to which they are attached denote a 6-membered heterocyclyl group, especially pyridyl.

Y may be, for example, a pyrimidinyl group of formula




where  $\text{R}^5$ ,  $\text{R}^6$  and  $\text{R}^7$  are each independently hydrogen,  $\text{C}_1\text{-C}_8$ -alkyl,  $\text{C}_1\text{-C}_8$ -alkoxy,  $\text{C}_1\text{-C}_8$ -alkylthio,  $\text{C}_1\text{-C}_8$ -alkylamino, di( $\text{C}_1\text{-C}_8$ -alkylamino or acylamino, or Y may be a group of formula




where  $\text{R}^8$ ,  $\text{R}^9$  and  $\text{R}^{10}$  are each independently hydrogen,  $\text{C}_1\text{-C}_8$ -alkyl,  $\text{C}_1\text{-C}_8$ -alkoxy,  $\text{C}_1\text{-C}_8$ -alkylthio,  $\text{C}_1\text{-C}_8$ -alkylamino, di( $\text{C}_1\text{-C}_8$ -alkyl) amino or acylamino.

Preferably Y is a group of formula



where  $\text{R}^5$  and  $\text{R}^6$  are each hydrogen and  $\text{R}^7$  is hydrogen,  $\text{C}_1\text{-C}_4$ -alkyl,  $\text{C}_1\text{-C}_4$ -alkylthio, or Y is a group of formula



where  $R^9$  and  $R^{10}$  are each hydrogen and  $R^8$  is hydrogen or di( $C_1$ - $C_4$ -alkyl) amino.

Preferred compounds of formula I in free or salt form are those where

Ar is phenyl optionally substituted by halogen or cyano,

$R^1$  is hydrogen, phenyl optionally substituted by cyano, halogen, carboxy or  $C_1$ - $C_4$ -alkoxy, or  $R^1$  is a monovalent 6-membered N-heterocyclic group,

$R^2$  is hydrogen,  $C_1$ - $C_4$ -alkylcarbonyl, 5-membered heterocyclylcarbonyl or phenylcarbonyl in which the phenyl moiety is optionally substituted by  $C_1$ - $C_8$ -alkoxy, and

Y is pyrimidinyl or pyridazinyl optionally substituted by  $C_1$ - $C_4$ -alkyl,  $C_1$ - $C_4$ -alkylthio,  $C_1$ - $C_4$ -alkylamino, di( $C_1$ - $C_4$ -alkyl) amino or  $C_1$ - $C_4$ -alkylcarbonylamino.

Further preferred compounds of formula I in free or salt form are those where

Ar is phenyl substituted by cyano meta to the indicated thiazole ring,

$R^1$  is hydrogen, phenyl substituted by cyano, fluorine, carboxy or  $C_1$ - $C_4$ -alkoxy or  $R^1$  is 6-membered N-heterocyclyl having one or two ring nitrogen atoms, optionally substituted by  $C_1$ - $C_4$ -alkyl or  $C_1$ - $C_4$ -alkoxy,

$R^2$  is hydrogen,  $C_1$ - $C_4$ -alkylcarbonyl, furylcarbonyl or  $C_1$ - $C_4$ -alkoxyphenylcarbonyl, and

Y is a group of formula IV or V as hereinbefore defined.

Especially preferred specific compounds of formula I are those described hereinafter in the Examples.

Many of the compounds represented by formula I are capable of forming acid addition salts, particularly pharmaceutically acceptable acid addition salts. Pharmaceutically acceptable acid addition salts of the compound of formula I include those of inorganic acids, for example, hydrohalic acids such as hydrofluoric acid, hydrochloric acid, hydrobromic acid or hydroiodic acid, nitric acid, sulfuric acid, phosphoric acid; and organic acids, for example aliphatic monocarboxylic acids such as formic acid, acetic acid, trifluoroacetic acid, propionic acid and butyric acid, aliphatic hydroxy acids such as lactic acid, citric acid, tartaric acid or malic acid, dicarboxylic acids such as maleic acid or succinic acid, aromatic carboxylic acids such as benzoic acid, p-chlorobenzoic acid, diphenylacetic acid or




triphenylacetic acid, aromatic hydroxy acids such as o-hydroxybenzoic acid, p-hydroxybenzoic acid, 1-hydroxynaphthalene-2-carboxylic acid or 3-hydroxynaphthalene-2-carboxylic acid, and sulfonic acids such as methanesulfonic acid or benzenesulfonic acid. These salts may be prepared from compounds of formula I by known salt-forming procedures.

Compounds of formula I which contain acidic, e.g. carboxyl, groups, are also capable of forming salts with bases, in particular pharmaceutically acceptable bases such as those well known in the art; suitable such salts include metal salts, particularly alkali metal or alkaline earth metal salts such as sodium, potassium, magnesium or calcium salts, or salts with ammonia or pharmaceutically acceptable organic amines or heterocyclic bases such as ethanolamines, benzylamines or pyridine. These salts may be prepared from compounds of formula I by known salt-forming procedures.


The invention provides, in another aspect, a method of preparing a compound of formula I in free or salt form which comprises

- (i) (A) for the preparation of compounds of formula I where  $R^1$  is optionally substituted phenyl or a 5- or 6- membered heterocyclic group, reacting a compound of formula



VI


in the form of a salt, e.g. a hydrohalide salt thereof, where Ar and Y are as hereinbefore defined and X is halogen, preferably bromine, with a compound of formula



VII

where  $R^1$  is phenyl optionally substituted by one or more substituents selected from halogen, cyano, hydroxy,  $\text{C}_1$ - $\text{C}_8$ -alkyl,  $\text{C}_1$ - $\text{C}_8$ -haloalkyl,  $\text{C}_1$ - $\text{C}_8$ -alkoxy,  $\text{C}_1$ - $\text{C}_8$ -alkoxy- $\text{C}_1$ - $\text{C}_8$ -alkyl and acyloxy or  $R^1$  is a 5- or 6- membered monovalent heterocyclic group, and  $R^2$  is H or  $\text{C}_1$ - $\text{C}_8$ -alkyl or

- (B) for the preparation of compounds of formula I where  $R^2$  is acyl or  $-\text{CON}(\text{R}^3)\text{R}^4$ , reacting a compound of formula



VIII


where Ar, R<sup>1</sup> and Y are as hereinbefore defined with, respectively, an acylating derivative of a carboxylic acid, for example the anhydride or acid chloride thereof, or with a compound of formula Cl-CON(R<sup>3</sup>)R<sup>4</sup>) where R<sup>3</sup> and R<sup>4</sup> are as hereinbefore defined, and

(ii) recovering the resultant compound of formula I in free or salt form.

Process variant (A) may be carried out in an organic solvent, for example an alcohol such as ethanol. Suitable reaction temperatures are elevated temperatures, for example from 50°C to reflux temperature of the solvent.

Process variant (B) may be carried out using known procedures for reaction of amines with acylating agents.


Compounds of formula VI may be prepared by reacting a compound of formula



IX

where Ar and Y are as hereinbefore defined, with halogen X<sub>2</sub>, preferably bromine. This halogenation may be effected using known procedures for alpha halogenation of ketones. Conveniently, the compound of formula VI is not isolated but is reacted directly with a compound of formula VII to give a compound of formula I.


Compounds of formula VII are thioureas which are either known or may be obtained by known procedures. For example they may be prepared by reaction of a compound of formula



X

where R<sup>1</sup> and R<sup>2</sup> are as hereinbefore defined, with benzoyl isothiocyanate and hydrolysing the resulting product, for example with aqueous NaOH, to replace the benzoyl group by halogen. The reaction with benzoyl isothiocyanate may be carried out in an organic solvent, for example an alcohol such as ethanol. Suitable reaction temperatures are from room temperature to reflux temperature of the solvent, conveniently 35-45°C. The hydrolysis may be effected at elevated temperature, for example 70°C to reflux temperature, conveniently at reflux temperature.

Pyrimidine, pyridazine and alkyl-substituted pyrimidine and pyridazine are known compounds which are commercially available or may be prepared by known procedures. Compounds of formula VIII may be prepared by process variant (A) as described above. Compounds of formula IX may be prepared by reaction of a compound of formula



with an alkali metal derivative of the pyrimidine or pyridazine, of formula MCH<sub>2</sub>Y where M is an alkali metal, preferably lithium or sodium, and Y is as hereinbefore defined, the CH<sub>2</sub>M group preferably being attached to the 4-position in the pyrimidinyl or pyridazinyl group, e.g. using a known procedure such as described hereinafter in the Examples. Compounds of formula X and XI are known or may be obtained by known procedures. For example, compounds of formula XI may be prepared by procedures such as described hereinafter in the Examples.

Compounds of formula I and their pharmaceutically acceptable salts are useful as pharmaceuticals. In particular, they exhibit inhibition of adenosine A2b receptor activation, i.e. they act as A2b receptor antagonists. Moreover, in general they selectively inhibit activation of A2b receptor over the adenosine A1 and A2a receptors. Their inhibitory properties may be demonstrated in the following test procedures:

#### Adenosine A2b Receptor Reporter Gene Assay

##### a) Culturing of Chinese Hamster Ovary (CHO) A2b Cell Line

CHO cells transfected with a Luciferase-expressing reporter plasmid (pCRE-LUCI) and with a plasmid carrying the human adenosine A2b receptor structural gene (pA2bRCV) are routinely cultured in Dulbecco's Modified Eagle Medium (DMEM) - supplemented with 10% v/v fetal calf serum (FCS), 2mM L-glutamine, 0.4 mg/ml L-proline, 1 nM sodium selenite, 0.5 mg/ml Hygromycin B and 1 mg/ml Geneticin - at 37°C, 5% CO<sub>2</sub> and 100% humidity. The cells are left to grow to confluence for 4-5 days. The cells obtained are passaged using trypsin/EDTA and split at a ratio of 1 in 5.

b) Preparation of cells for assay

Prior to the assay, the CHO-A2b cells are plated onto white 96-well View Plate tissue culture plates (Packard) at a density of 50,000 cells per well in 50 µl of DMEM, and the plates are incubated at 37°C, 5% CO<sub>2</sub> and 100 % humidity.

c) Preparation of Reference and Test Compounds

10 mM solutions of the reference compound, Xanthine Amine Cogener (XAC), and the test compound in dimethyl sulfoxide (DMSO) are prepared. The solutions are further diluted with DMSO to 100 µM, then diluted to 10 µM, and finally to 250 nM or 2.5 µM with Assay Buffer (DMEM Phenol Red-free tissue culture media supplemented with 10 µM Rolipram and 10 U/ml adenosine deaminase (ADA). The resulting solutions (40 µl) are added to the cells in the appropriate wells, the final concentration per well being 100 nM or 1 µM, and the plates are incubated at 37°C, 5% CO<sub>2</sub> and 100% humidity.

d) Luciferase Reporter Gene Assay

5'-N-ethylcarboxamidoadenosine (NECA), an adenosine A2b agonist, is prepared as a 10 nM solution in DMSO and then diluted to 100 µM with Assay Buffer. This solution is serially diluted in Assay Buffer to give a series of 10 NECA concentrations from 100 to 0.01 µM. 10 µl portions of the resulting NECA solutions are added to the mixtures of CHO-A2b cells and reference or test compound solutions prepared as described above (preincubated for 30 minutes), final concentrations ranging from 10 to 0.0005 µM per well. The cells are incubated at 37°C, 5% CO<sub>2</sub> and 100% humidity for 3 hours to induce release of cAMP, which then binds to cAMP binding protein (CBP) and the resulting complex interacts with

the reporter plasmid to express Luciferase. 100  $\mu$ l of Steady-Glo, a Luciferase assay substrate from Promega, is added to all wells to lyse the cells and generate luminescence in proportion to the amount of Luciferase produced. The plates are left for a minimum of 5 minutes before being read on the luminescence program of a Topcount NXT microplate scintillation counter (ex Packard). Concentration - response curves are plotted from the luminescence data using Activitybase software and  $K_B$  values for the antagonists under test are calculated from the shifts of the curve at a particular concentration ( $K_B = [\text{antagonist}]/(\text{concentration ratio} - 1)$ )

Compounds of the Examples hereinbelow have  $K_B$  values below 100nM in the reporter gene assay. For example, the compounds of Examples 4,6,9,12 and 16 have  $K_B$  values of 38, 29, 6, 2 and 50nM respectively.

Having regard to their inhibition of adenosine A2b receptor activation, compounds of formula I in free or pharmaceutically acceptable salt form, hereinafter alternately referred to as agents of the invention, are useful in the treatment of conditions which are mediated by the activation of the adenosine A2b receptor, particularly inflammatory or allergic conditions. Treatment in accordance with the invention may be symptomatic or prophylactic.

Accordingly, agents of the invention are useful in the treatment of inflammatory or obstructive airways diseases, resulting, for example, in reduction of tissue damage, bronchial hyperreactivity, remodelling or disease progression. Inflammatory or obstructive airways diseases to which the present invention is applicable include asthma of whatever type or genesis including both intrinsic (non-allergic) asthma and extrinsic (allergic) asthma, mild asthma, moderate asthma, severe asthma, bronchitic asthma, exercise-induced asthma, occupational asthma and asthma induced following bacterial infection. Treatment of asthma is also to be understood as embracing treatment of subjects, e.g. of less than 4 or 5 years of age, exhibiting wheezing symptoms and diagnosed or diagnosable as "wheezy infants", an established patient category of major medical concern and now often identified as incipient or early-phase asthmatics. (For convenience this particular asthmatic condition is referred to as "wheezy-infant syndrome".)

Prophylactic efficacy in the treatment of asthma will be evidenced by reduced frequency or severity of symptomatic attack, e.g. of acute asthmatic or bronchoconstrictor attack,

improvement in lung function or improved airways hyperreactivity. It may further be evidenced by reduced requirement for other, symptomatic therapy, i.e. therapy for or intended to restrict or abort symptomatic attack when it occurs, for example anti-inflammatory (e.g. corticosteroid) or bronchodilatory. Prophylactic benefit in asthma may in particular be apparent in subjects prone to "morning dipping". "Morning dipping" is a recognised asthmatic syndrome, common to a substantial percentage of asthmatics and characterised by asthma attack, e.g. between the hours of about 4 to 6 am, i.e. at a time normally substantially distant from any previously administered symptomatic asthma therapy.

Other inflammatory or obstructive airways diseases and conditions to which the present invention is applicable include acute lung injury (ALI), adult respiratory distress syndrome (ARDS), chronic obstructive pulmonary, airways or lung disease (COPD, COAD or COLD), including chronic bronchitis or dyspnea associated therewith, emphysema, as well as exacerbation of airways hyperreactivity consequent to other drug therapy, in particular other inhaled drug therapy. The invention is also applicable to the treatment of bronchitis of whatever type or genesis including, e.g., acute, arachidic, catarrhal, croupus, chronic or phthinoïd bronchitis. Further inflammatory or obstructive airways diseases to which the present invention is applicable include pneumoconiosis (an inflammatory, commonly occupational, disease of the lungs, frequently accompanied by airways obstruction, whether chronic or acute, and occasioned by repeated inhalation of dusts) of whatever type or genesis, including, for example, aluminosis, anthracosis, asbestosis, chalicosis, ptilosis, siderosis, silicosis, tabacosis and byssinosis.

Having regard to their anti-inflammatory activity, in particular in relation to inhibition of eosinophil activation, agents of the invention are also useful in the treatment of eosinophil related disorders, e.g. eosinophilia, in particular eosinophil related disorders of the airways (e.g. involving morbid eosinophilic infiltration of pulmonary tissues) including hypereosinophilia as it effects the airways and/or lungs as well as, for example, eosinophil-related disorders of the airways consequential or concomitant to Löffler's syndrome, eosinophilic pneumonia, parasitic (in particular metazoan) infestation (including tropical eosinophilia), bronchopulmonary aspergillosis, polyarteritis nodosa (including Churg-Strauss syndrome), eosinophilic granuloma and eosinophil-related disorders affecting the airways occasioned by drug-reaction.

Agents of the invention are also useful in the treatment of inflammatory or allergic conditions of the skin, for example psoriasis, contact dermatitis, atopic dermatitis, alopecia areata, erythema multiforma, dermatitis herpetiformis, scleroderma, vitiligo, hypersensitivity angitis, urticaria, bullous pemphigoid, lupus erythematosus, pemphigus, epidermolysis bullosa acquisita, and other inflammatory or allergic conditions of the skin.

Agents of the invention may also be used for the treatment of other diseases or conditions, in particular diseases or conditions having an inflammatory component, for example, treatment of diseases and conditions of the eye such as conjunctivitis, keratoconjunctivitis sicca, and vernal conjunctivitis, diseases affecting the nose including allergic rhinitis, and inflammatory disease in which autoimmune reactions are implicated or having an autoimmune component or aetiology, including autoimmune haematological disorders (e.g. haemolytic anaemia, aplastic anaemia, pure red cell anaemia and idiopathic thrombocytopenia), systemic lupus erythematosus, polychondritis, scleroderma, Wegener granulomatosis, dermatomyositis, chronic active hepatitis, myasthenia gravis, Steven-Johnson syndrome, idiopathic sprue, autoimmune inflammatory bowel disease (e.g. ulcerative colitis and Crohn's disease), endocrine ophthalmopathy, Grave's disease, sarcoidosis, alveolitis, chronic hypersensitivity pneumonitis, multiple sclerosis, primary biliary cirrhosis, diabetes, juvenile diabetes (diabetes mellitus type I), uveitis (anterior and posterior), keratoconjunctivitis sicca and vernal keratoconjunctivitis, interstitial lung fibrosis, psoriatic arthritis and glomerulonephritis (with and without nephrotic syndrome, e.g. including idiopathic nephrotic syndrome or minimal change nephropathy).

Other diseases or conditions mediated by the adenosine A2b receptor which may be treated with agents of the invention include diarrheal diseases, ischemia/reperfusion injuries or retinopathy such as diabetic retinopathy or hyperbaric oxygen-induced retinopathy.

In accordance with the foregoing, the invention also provides a method for the treatment of a condition mediated by activation of the adenosine A2b receptor, for example an inflammatory or allergic condition, particularly an inflammatory or obstructive airways disease, which comprises administering to a subject, particularly a human subject, in need thereof an effective amount of a compound of formula I in free form or in the form of a pharmaceutically acceptable salt. In another aspect the invention provides a compound of formula I, in free form or in the form of a pharmaceutically acceptable salt, for use in the

manufacture of a medicament for the treatment of a condition mediated by activation of the adenosine A2b receptor, particularly an inflammatory or obstructive airways disease.

The agents of the invention may be administered by any appropriate route, e.g. orally, for example in the form of a tablet or capsule; parenterally, for example intravenously; by inhalation, for example in the treatment of inflammatory or obstructive airways disease; intranasally, for example in the treatment of allergic rhinitis; topically to the skin, for example in the treatment of atopic dermatitis; or rectally, for example in the treatment of inflammatory bowel disease.

In a further aspect, the invention also provides a pharmaceutical composition comprising a compound of formula I in free form or in the form of a pharmaceutically acceptable salt, optionally together with a pharmaceutically acceptable diluent or carrier therefor. Such compositions may be prepared using conventional diluents or excipients and techniques known in the galenic art. Thus oral dosage forms may include tablets and capsules. Formulations for topical administration may take the form of creams, ointments, gels or transdermal delivery systems, e.g. patches. Compositions for inhalation may comprise aerosol or other atomizable formulations or dry powder formulations.


Dosages of agents of the invention employed in practising the present invention will of course vary depending, for example, on the particular condition to be treated, the effect desired and the mode of administration. In general, suitable daily dosages for oral administration are of the order of 0.1 to 10 mg/kg.

The invention is illustrated by the following Examples.

#### Examples 1-12







Compounds of formula I which are also of formula






are shown in the following table, the method of preparation being described hereinafter. The table also shows mass spectrometry data. The Examples are in free form.

Ex. No.	R <sup>a</sup>	R <sup>b</sup>	R <sup>c</sup>	R <sup>1</sup>	R <sup>2</sup>	m/s (MH <sup>+</sup> )
1	CN	H	CH <sub>3</sub>		H	371.2
2	CN	H	H		H	381
3	CN	H	H		H	385.5
4	CN	H	H		H	371.4
5	CN	H	CH <sub>3</sub>		H	371.2
6	CN	H	H		H	381




7	CN	H	H		H	371
8	CN	H	H		H	387.5
9	CN	H	H		H	357.1
10	CN	H	H	H		374.1
11	CN	H	CH <sub>3</sub>	H		414
12	CN	H	H		H	358

### Examples 13-16

Compounds of formula I which are also of formula



are shown in the following table, the method of preparation being described hereinafter. The table also shows mass spectrometry data. The Examples are in free form, with the exception of Example 16 which is in the form of a salt with trifluoroacetic acid.

Ex.	R <sup>a</sup>	R <sup>b</sup>	R <sup>d</sup>	R <sup>1</sup>	R <sup>2</sup>	M/s MH <sup>+</sup>
13	CN	H	H	H	COCH <sub>3</sub>	322.4
14	CN	H	H		H	357.4
15	CN	H	H		H	400.4
16	CN	H	N(CH <sub>3</sub> ) <sub>2</sub>		H	417.5

#### Preparation of 3-cyano-N-methoxy-N-methylbenzamide

3-cyanobenzoic acid (7.00g, 47.6mmol) is dissolved in THF (50ml) under nitrogen and N,N'-carbonylbisimidazole (7.95g, 49.0mmol) is added. The mixture is stirred under reflux for 25mins, then cooled and dimethylhydroxylamine hydrochloride (4.78g, 49.0mmol) is added. The mixture is heated under reflux for 5 hours and then stood at room temperature for 18h. The reaction mixture is added to water (200ml). The aqueous mixture is extracted with ether twice, and the combined ether phases are washed with saturated aqueous sodium carbonate. The organic layer is dried (MgSO<sub>4</sub>) and concentrated to a colourless mobile oil.

MS: (APCI) MH<sup>+</sup> at 191

#### Preparation of compounds of formula IX

##### 3-(Pyridazin-4-yl-acetyl)-benzonitrile

A solution of diisopropylamine (4.08ml, 29.0mmol) in dry THF (35ml) is cooled to -70°C and 1.6M BuLi in hexanes (18.1ml, 29.0mmol) is added dropwise. The cooling bath is then

removed and the mixture is allowed to warm to 0°C. The solution is recooled to -70°C and 4-methylpyridazine (2.5g, 26.6mmol) is added dropwise, and the solution stirred at -70°C for 1.75h. To the resulting brown solution 3-cyano-N-methoxy-N-methyl-benzamide (5.06g, 26.6mmol) in dry tetrahydrofuran (10ml) is added dropwise. The solution is stirred at -70°C for 1h, then warmed to -30°C. After 1h the mixture is allowed to warm to room temperature. Saturated aqueous ammonium chloride (20ml) is added to the reaction, and the solvent is removed by evaporation to yield a yellow solid, which is washed with aqueous sodium bicarbonate, water and then ethyl acetate. The product is obtained as a yellow solid, m.p. 142-144°C. MS (MH<sup>+</sup>) 224

### 3-(Pyrimidin-4-yl-acetyl)-benzonitrile

A solution of diisopropylamine (4.1ml, 29.3mmol) in dry THF (40ml) is cooled to -60°C and 1.6M BuLi in hexanes (16.6ml, 26.6mmol) is added dropwise. The mixture is then allowed to warm to 20°C. The solution is recooled to -60°C, 4-methylpyrimidine (2.5g, 26.6mmol) is added dropwise, and the solution is stirred at -60°C for 1h. To the resulting solution 3-cyano-N-methoxy-N-methyl-benzamide (5.05g, 26.6mmol) in dry tetrahydrofuran (10ml) is added dropwise. The solution is stirred at -60°C for 1h, then allowed to warm to room temperature. After 18h, water is added to the reaction mixture, and the solvent is removed by evaporation to yield a solid. Water is added to the mass, and the resulting solid filtered off and washed with water. The product is obtained as a solid, MS (APCI, MH<sup>+</sup>) 224.

The following compounds are prepared in an analogous manner:

3-[(6-Methyl-pyrimidin-4-yl)-acetyl]-benzonitrile, MS (MH<sup>+</sup>) 238.1

3-[(6-Dimethylamino-pyridazin-4-yl)-acetyl]-benzonitrile, MS (MH<sup>+</sup>) 267.3

### Preparation of compounds of formula VIII

#### 3-(2-Amino-5-pyrimidin-4-yl-thiazol-4-yl)-benzonitrile

3-(Pyrimidin-4-yl-acetyl)-benzonitrile (1.00g, 4.48mmol) is dissolved in dry dioxane (20ml) and bromine (0.23ml, 4.48mmol) is added dropwise. The mixture is stirred for 30 minutes and then the solvent is evaporated under vacuum at 30°C to give a gum. This residue is dissolved in absolute ethanol (11.5ml). To 7ml of this solution thiourea (0.188g, 2.17mmol) is added and the mixture is heated at a gentle reflux for 6 hours. After cooling the reaction mixture, the solid is filtered off and washed with ethanol. The solid is then suspended in

water and aqueous ammonia added dropwise. The resulting solid is filtered off, washed with water and dried. MS (MH<sup>+</sup>) 280.1

The following compounds are prepared in an analogous manner:

3-[2-Amino-5-(6-methyl-pyrimidin-4-yl)-thiazol-4-yl]-benzonitrile; MS (MH<sup>+</sup>) 294.4

3-(2-Amino-5-pyridazin-4-yl-thiazol-4-yl)-benzonitrile, MS (MH<sup>+</sup>) 280.3

#### Preparation of Specific Examples

N-[4-(3-Cyano-phenyl)-5-pyrimidin-4-yl-thiazol-2-yl]-3-methoxy-benzamide (Example 11)

3-(2-Amino-5-pyrimidin-4-yl-thiazol-4-yl)-benzonitrile (90mg, 0.32mmol) is suspended in pyridine (0.75ml) and 3-methoxybenzoyl chloride (165mg, 0.96mmol) is added. A precipitate is formed. The reaction mixture is stirred at room temperature for 18 hours. An excess of water is added to the reaction mixture, and the resulting solid is collected, washed with water and dried. The dried solid is triturated with hot ethanol to give product as a pale powder. m.p. 255-256°C, MS (APCI MH<sup>+</sup>) 414. Examples 10 and 13 are prepared analogously.

3-[2-(Pyrazin-2-ylamino)-5-pyrimidin-4-yl-thiazol-4-yl]-benzonitrile (Example 12)

3-(pyrimidin-4-yl-acetyl)-benzonitrile (1.00g, 4.48mmol) is dissolved in dry dioxane (11ml) and bromine (0.23ml, 4.48mmol) is added dropwise. The mixture is stirred for 30 minutes and then the solvent is evaporated under vacuum at 30°C to give a solid. This residue is dissolved in absolute ethanol (10.5ml). To 1.5ml of this solution pyrazin-2-yl-thiourea (108mg, 0.70mmol) is then added and the mixture is heated at 70°C for 16 hours. The crude reaction mixture is diluted with water and the resulting solid collected and washed with ethanol. This solid is suspended in water and aqueous ammonia is added to pH 9. After 10 min. the resulting solid is collected by filtration and washed with ethanol/ethyl acetate mixtures. The solid is dried under vacuum at 70°C to give a powder. m.p. >280°C MS (APCI) MH<sup>+</sup> and, MH<sup>-</sup> 358 and 356 respectively. Examples 1 to 9 are prepared analogously.


4-[4-(3-Cyano-phenyl)-5-pyridazin-4-yl-thiazol-2-ylamino]-benzoic acid (Example 15)

3-(pyridazin-4-yl-acetyl)-benzonitrile (1.55g, 6.94mmol) is dissolved in dry dioxane (20ml) at 10-15°C and bromine (0.35ml, 6.9mmol) is added dropwise. The mixture is stirred for

45 minutes and the resulting solid is dissolved in methanol and the solution evaporated to a gum under vacuum at room temperature. This residue is dissolved in dry dimethylformamide (12ml). To 4ml of this solution 4-carboxyphenyl thiourea (0.496g, 2.5mmol) is then added and the mixture is heated at 70°C for 20 hours. The reaction mixture is allowed to cool and the solvent removed under vacuum. The residue is triturated with ethyl acetate and then ammonia solution is added. The resulting solid is filtered off, washed with water and dried to give a solid, m.p. >275°C. MS (MH<sup>+</sup>) 400.4. Examples 14 and 16 are prepared analogously.

Claims

1. A compound of formula



in free or salt form, where

Ar is a C<sub>6</sub>-C<sub>15</sub> monovalent aromatic group,

R<sup>1</sup> is hydrogen, phenyl optionally substituted by one or more substituents selected from halogen, cyano, hydroxy, C<sub>1</sub>-C<sub>8</sub>-alkyl, C<sub>1</sub>-C<sub>8</sub>-haloalkyl, C<sub>1</sub>-C<sub>8</sub>-alkoxy, C<sub>1</sub>-C<sub>8</sub>-alkoxy-C<sub>1</sub>-C<sub>8</sub>-alkyl and acyloxy, or R<sup>1</sup> is a 5- or 6- membered monovalent heterocyclic group,

R<sup>2</sup> is hydrogen, C<sub>1</sub>-C<sub>8</sub>-alkyl, acyl or -CON(R<sup>3</sup>)R<sup>4</sup>,

R<sup>3</sup> and R<sup>4</sup> are each independently hydrogen or C<sub>1</sub>-C<sub>8</sub>-alkyl, or together with the nitrogen atom to which they are attached denote a 5- or 6- membered heterocyclic group, and


Y is a pyrimidinyl or pyridazinyl group, optionally substituted by at least one C<sub>1</sub>-C<sub>8</sub>-alkyl, C<sub>1</sub>-C<sub>8</sub>-alkoxy, C<sub>1</sub>-C<sub>8</sub>-alkylthio, C<sub>1</sub>-C<sub>8</sub>-alkylamino, di(C<sub>1</sub>-C<sub>8</sub>-alkyl)amino or acylamino group.

2. A compound according to claim 1, in which Ar is phenyl optionally substituted by halogen, cyano or C<sub>1</sub>-C<sub>8</sub>-alkyl.


3. A compound according to claim 1 or 2, in which R<sup>1</sup> is phenyl optionally substituted by cyano, carboxy or C<sub>1</sub>-C<sub>4</sub>-alkoxy, or R<sup>1</sup> is a monovalent 6-membered N-heterocyclic group.

4. A compound according to claim 1, 2 or 3, in which R<sup>2</sup> is hydrogen, C<sub>1</sub>-C<sub>4</sub>-alkylcarbonyl, 5-membered heterocyclylcarbonyl, or phenylcarbonyl in which the phenyl moiety is optionally substituted by C<sub>1</sub>-C<sub>8</sub>-alkoxy.

5. A compound according to one of claims 1 to 4, in which Y is a group of formula



where R<sup>5</sup> and R<sup>6</sup> are each hydrogen and R<sup>7</sup> is hydrogen, C<sub>1</sub>-C<sub>4</sub>-alkyl or C<sub>1</sub>-C<sub>4</sub>-alkylthio, or Y is a group of formula



where R<sup>9</sup> and R<sup>10</sup> are each hydrogen and R<sup>8</sup> is hydrogen or di(C<sub>1</sub>-C<sub>4</sub>-alkyl)amino.

6. A compound according to claim 1, in which

Ar is phenyl optionally substituted by halogen or cyano,

R<sup>1</sup> is hydrogen, phenyl optionally substituted by cyano, halogen, carboxy or C<sub>1</sub>-C<sub>4</sub>-alkoxy, or R<sup>1</sup> is a monovalent 6-membered N-heterocyclic group,

R<sup>2</sup> is hydrogen, C<sub>1</sub>-C<sub>4</sub>-alkylcarbonyl, 5-membered heterocyclylcarbonyl or phenylcarbonyl in which the phenyl moiety is optionally substituted by C<sub>1</sub>-C<sub>8</sub>-alkoxy, and

Y is pyrimidinyl or pyridazinyl optionally substituted by C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy, C<sub>1</sub>-C<sub>4</sub>-alkylthio, C<sub>1</sub>-C<sub>4</sub>-alkylamino, di(C<sub>1</sub>-C<sub>4</sub>-alkyl) amino or C<sub>1</sub>-C<sub>4</sub>-alkylcarbonylamino.

7. A compound according to claim 1, in which

Ar is phenyl substituted by cyano meta to the indicated thiazole ring,

R<sup>1</sup> is hydrogen, phenyl substituted by cyano, fluorine, carboxy or C<sub>1</sub>-C<sub>4</sub>-alkoxy or R<sup>1</sup> is 6-membered N-heterocyclyl having one or two ring nitrogen atoms, optionally substituted by C<sub>1</sub>-C<sub>4</sub>-alkyl or C<sub>1</sub>-C<sub>4</sub>-alkoxy,

R<sup>2</sup> is hydrogen, C<sub>1</sub>-C<sub>4</sub>-alkylcarbonyl, furylcarbonyl or C<sub>1</sub>-C<sub>4</sub>-alkoxyphenylcarbonyl, and

Y is a group of formula IV or V as defined in claim 5.

8. A compound according to any one of claims 1 to 7 for use as a pharmaceutical.




9. A pharmaceutical composition comprising a compound according to any one of claims 1 to 7, optionally together with a pharmaceutically acceptable diluent or carrier.


10. The use of a compound according to any one of claims 1 to 7 in the manufacture of a medicament for the treatment of a condition mediated by activation of the adenosine A2b receptor.

11. The use of a compound according to any one of claims 1 to 7 in the manufacture of a medicament for the treatment of an inflammatory or obstructive airways disease.

12. A method of preparing a compound of formula I in free or salt form which comprises  
(i) (A) for the preparation of compounds of formula I where  $R^1$  is optionally substituted phenyl or a 5- or 6- membered heterocyclic group, reacting a compound of formula




in the form of a salt, where Ar and Y are as defined in claim 1 and X is halogen, with a compound of formula



where  $R^1$  is phenyl optionally substituted by one or more substituents selected from halogen, cyano, hydroxy,  $C_1$ - $C_8$ -alkyl,  $C_1$ - $C_8$ -haloalkyl,  $C_1$ - $C_8$ -alkoxy,  $C_1$ - $C_8$ -alkoxy- $C_1$ - $C_8$ -alkyl and acyloxy or  $R^1$  is a 5- or 6- membered monovalent heterocyclic group, and  $R^2$  is H or  $C_1$ - $C_8$ -alkyl or

(B) for the preparation of compounds of formula I where  $R^2$  is acyl or  $-CON(R^3)R^4$ , reacting a compound of formula



where Ar, R<sup>1</sup> and Y are as hereinbefore defined with, respectively, an acylating derivative of a carboxylic acid or with a compound of formula Cl-CON(R<sup>3</sup>)R<sup>4</sup>) where R<sup>3</sup> and R<sup>4</sup> are as defined in claim 1, and

(ii) recovering the resultant compound of formula I in free or salt form.